

More Examples of Pseudo-Collars on High-Dimensional Manifolds

Dr. Jeffrey Rolland

Department of Mathematics
Milwaukee School of Engineering

February 03, 2020

Quillen's Plus Construction

- **Quillen's plus construction** is a way of taking a CW complex or manifold X whose fundamental group G contains a perfect normal subgroup P and creating a new CW complex or manifold X^+ whose fundamental group is simpler, namely $Q = G/P$

Quillen's Plus Construction

- **Quillen's plus construction** is a way of taking a CW complex or manifold X whose fundamental group G contains a perfect normal subgroup P and creating a new CW complex or manifold X^+ whose fundamental group is simpler, namely $Q = G/P$
- If the object is a manifold, the plus construction creates a **plus cobordism** (W, X, X^+) between the two manifold with $X^+ \hookrightarrow W$ a simple homotopy equivalence which gives rise to a map f between the two manifolds

Quillen's Plus Construction

- **Quillen's plus construction** is a way of taking a CW complex or manifold X whose fundamental group G contains a perfect normal subgroup P and creating a new CW complex or manifold X^+ whose fundamental group is simpler, namely $Q = G/P$
- If the object is a manifold, the plus construction creates a **plus cobordism** (W, X, X^+) between the two manifold with $X^+ \hookrightarrow W$ a simple homotopy equivalence which gives rise to a map f between the two manifolds
- If the object is a CW complex, the Plus Construction simply create a map f between the two CW complexes

Quillen's Plus Construction

- **Quillen's plus construction** is a way of taking a CW complex or manifold X whose fundamental group G contains a perfect normal subgroup P and creating a new CW complex or manifold X^+ whose fundamental group is simpler, namely $Q = G/P$
- If the object is a manifold, the plus construction creates a **plus cobordism** (W, X, X^+) between the two manifold with $X^+ \hookrightarrow W$ a simple homotopy equivalence which gives rise to a map f between the two manifolds
- If the object is a CW complex, the Plus Construction simply create a map f between the two CW complexes
- In either case, the map f is a $\mathbb{Z}Q$ -homology isomorphism

Semi-S-Cobordisms

- In previous work, we had discovered a partial reverse to Quillen's plus construction in the high-dimensional manifold category

Semi-S-Cobordisms

- In previous work, we had discovered a partial reverse to Quillen's plus construction in the high-dimensional manifold category

Theorem 1 (R., 2015)

Let M be a manifold of dimension 6 or higher, with $\pi_1(M) \cong Q$. Let K be a finitely presented superperfect group. Let G be a semi-direct product of Q by K , $G = K \rtimes Q$. Then there is a cobordism (W, M, M_-) with $\pi_1(M_-) \cong G$ and $M \hookrightarrow W$ a simple homotopy equivalence

- In previous work, we had discovered a partial reverse to Quillen's plus construction in the high-dimensional manifold category

Theorem 1 (R., 2015)

Let M be a manifold of dimension 6 or higher, with $\pi_1(M) \cong Q$. Let K be a finitely presented superperfect group. Let G be a semi-direct product of Q by K , $G = K \rtimes Q$. Then there is a cobordism (W, M, M_-) with $\pi_1(M_-) \cong G$ and $M \hookrightarrow W$ a simple homotopy equivalence

- (Note that G being a semi-direct product of Q by K , $G = K \rtimes Q$, means G satisfies $1 \rightarrow K \rightarrow G \rightarrow Q \rightarrow 1$, so G is a group extension of Q by K , with a special condition for how elements of Q multiply elements of K in G)

Semi-S-Cobordisms

- In previous work, we had discovered a partial reverse to Quillen's plus construction in the high-dimensional manifold category

Theorem 1 (R., 2015)

Let M be a manifold of dimension 6 or higher, with $\pi_1(M) \cong Q$. Let K be a finitely presented superperfect group. Let G be a semi-direct product of Q by K , $G = K \rtimes Q$. Then there is a cobordism (W, M, M_-) with $\pi_1(M_-) \cong G$ and $M \hookrightarrow W$ a simple homotopy equivalence

- (Note that G being a semi-direct product of Q by K , $G = K \rtimes Q$, means G satisfies $1 \rightarrow K \rightarrow G \rightarrow Q \rightarrow 1$, so G is a group extension of Q by K , with a special condition for how elements of Q multiply elements of K in G)
- (Semi-direct products are the simplest kind of group extensions; direct products are one example)

- We call (W, M, M_-) a **semi-s-cobordism**, because it is “half an s-cobordism”

Semi-S-Cobordisms

- We call (W, M, M_-) a **semi-s-cobordism**, because it is “half an s-cobordism”
- Note (W, M_-, M) (read upside-down, with the roles of M and M_- reversed) is a plus cobordism (so $(M_-)^+ \approx M$)

Semi-S-Cobordisms

- What we would like to do now is “stack” these semi-s-cobordisms, forming (W_1, M, M_-) , (W_2, M_-, M_{--}) , and so on, out to infinity

- What we would like to do now is “stack” these semi-s-cobordisms, forming (W_1, M, M_-) , (W_2, M_-, M_{--}) , and so on, out to infinity
- Gluing W_1 and W_2 together across M_- and so on produces an $(n + 1)$ -dimensional, 1-ended manifold V whose neighborhoods of infinity are **pseudo-collared**

Semi-S-Cobordisms

- Pseudo-collars are generalizations of collar structures on the boundary of a manifold

Semi-S-Cobordisms

- Pseudo-collars are generalizations of collar structures on the boundary of a manifold
- The notion of a pseudo-collar originated in Hilbert cube manifold theory, where they were a necessary and sufficient condition for finding a \mathcal{Z} -set as the boundary of a manifold

Semi-S-Cobordisms

- Pseudo-collar are generalizations of collar structures on the boundary of a manifold
- The notion of a pseudo-collar originated in Hilbert cube manifold theory, where they were a necessary and sufficient condition for finding a \mathcal{Z} -set as the boundary of a manifold
- We would like to produce many examples of open, 1-ended, $(n + 1)$ -dimensional pseudo-collared manifolds

Semi-S-Cobordisms

- Pseudo-collar are generalizations of collar structures on the boundary of a manifold
- The notion of a pseudo-collar originated in Hilbert cube manifold theory, where they were a necessary and sufficient condition for finding a \mathcal{Z} -set as the boundary of a manifold
- We would like to produce many examples of open, 1-ended, $(n + 1)$ -dimensional pseudo-collared manifolds
- We could just keep changing the kernel group K at each stage, but we would like to keep the same kernel group K

Semi-S-Cobordisms

- Pseudo-collars are generalizations of collar structures on the boundary of a manifold
- The notion of a pseudo-collar originated in Hilbert cube manifold theory, where they were a necessary and sufficient condition for finding a \mathcal{Z} -set as the boundary of a manifold
- We would like to produce many examples of open, 1-ended, $(n + 1)$ -dimensional pseudo-collared manifolds
- We could just keep changing the kernel group K at each stage, but we would like to keep the same kernel group K
- The idea is that there will be infinitely many outer automorphisms, so we can form infinitely many semi-direct products, each with a different outer automorphism

Semi-S-Cobordisms

- Pseudo-collars are generalizations of collar structures on the boundary of a manifold
- The notion of a pseudo-collar originated in Hilbert cube manifold theory, where they were a necessary and sufficient condition for finding a \mathcal{Z} -set as the boundary of a manifold
- We would like to produce many examples of open, 1-ended, $(n + 1)$ -dimensional pseudo-collared manifolds
- We could just keep changing the kernel group K at each stage, but we would like to keep the same kernel group K
- The idea is that there will be infinitely many outer automorphisms, so we can form infinitely many semi-direct products, each with a different outer automorphism
- This is one advantage of using semi-direct products over direct products

Uncountable Many Pseudo-Collars

- A subset A of a finitely presented group S is called an unpermutable set if it is countably infinite and has the property that if $\phi : K \rightarrow K$ is an isomorphism and $\phi(a_i) = a_j^{\pm 1}$, then $i = j$

Uncountable Many Pseudo-Collars

- A subset A of a finitely presented group S is called an unpermutable set if it is countably infinite and has the property that if $\phi : K \rightarrow K$ is an isomorphism and $\phi(a_i) = a_j^{\pm 1}$, then $i = j$
- A large source of examples of unpermutable sets are fundamental groups of hyperbolic manifolds

Uncountable Many Pseudo-Collars

- A subset A of a finitely presented group S is called an unpermutable set if it is countably infinite and has the property that if $\phi : K \rightarrow K$ is an isomorphism and $\phi(a_i) = a_j^{\pm 1}$, then $i = j$
- A large source of examples of unpermutable sets are fundamental groups of hyperbolic manifolds
- If you take a geodesic loop γ that's non-null-homotopic, then the length of this loop is an invariant of this loop by Mostow Rigidity, so $A = \{\gamma^n \mid n \in \mathbb{N}\}$ forms an unpermutable set.

Uncountable Many Pseudo-Collars

- A subset A of a finitely presented group S is called an unpermutable set if it is countably infinite and has the property that if $\phi : K \rightarrow K$ is an isomorphism and $\phi(a_i) = a_j^{\pm 1}$, then $i = j$
- A large source of examples of unpermutable sets are fundamental groups of hyperbolic manifolds
- If you take a geodesic loop γ that's non-null-homotopic, then the length of this loop is an invariant of this loop by Mostow Rigidity, so $A = \{\gamma^n \mid n \in \mathbb{N}\}$ forms an unpermutable set.

Theorem 2 (R., 2020)

*Let M be a manifold of dimension $n \geq 6$ with fundamental group \mathbb{Z} . Let S be a finitely presented, superperfect, centerless, freely indecomposable, Hopfian, co-Hopfian group with an unpermutable set, and let $K = S * S$. Then there are uncountably many $(n + 1)$ -dimensional, pseudo-collarable, 1-ended manifolds V with boundary M*

- V will break up into semi-s-cobordisms (W_j, M_{j-1}, M_j) , where $G_j \cong S \times G_{j-1}$, $G_j = \pi_1(M_j)$

- V will break up into semi-s-cobordisms (W_j, M_{j-1}, M_j) , where $G_j \cong S \rtimes G_{j-1}$, $G_j = \pi_1(M_j)$
- Each G_j will, in fact, be isomorphic to $(S \times S \times \dots \times S) \rtimes \mathbb{Z}$

- V will break up into semi-s-cobordisms (W_j, M_{j-1}, M_j) , where $G_j \cong S \rtimes G_{j-1}$, $G_j = \pi_1(M_j)$
- Each G_j will, in fact, be isomorphic to $(S \times S \times \dots \times S) \rtimes \mathbb{Z}$
- We will produce different G_j 's by varying the outer actions, a technical part of semi-direct products, while keeping the quotient group, \mathbb{Z} , and kernel group, S , essentially constant

- V will break up into semi-s-cobordisms (W_j, M_{j-1}, M_j) , where $G_j \cong S \rtimes G_{j-1}$, $G_j = \pi_1(M_j)$
- Each G_j will, in fact, be isomorphic to $(S \times S \times \dots \times S) \rtimes \mathbb{Z}$
- We will produce different G_j 's by varying the outer actions, a technical part of semi-direct products, while keeping the quotient group, \mathbb{Z} , and kernel group, S , essentially constant
- We will produce one V for each $\omega \in \prod_{i=1}^{\infty} \{0, 1\}$

Uncountable Many Pseudo-Collars

- Of course, it's pretty easy to whip out a 1-ended, pseudo-collarable manifold V : you just keep taking larger and larger G_j 's and M_j 's and glue the cobordisms together

Uncountable Many Pseudo-Collars

- Of course, it's pretty easy to whip out a 1-ended, pseudo-collarable manifold V : you just keep taking larger and larger G_j 's and M_j 's and glue the cobordisms together
- The hard part is proving that the resulting pro-fundamental group systems at infinity are all non-isomorphic

Uncountable Many Pseudo-Collars

- Of course, it's pretty easy to whip out a 1-ended, pseudo-collarable manifold V : you just keep taking larger and larger G_j 's and M_j 's and glue the cobordisms together
- The hard part is proving that the resulting pro-fundamental group systems at infinity are all non-isomorphic
- For example, if $Q = \prod_{i=1}^{\infty} \mathbb{Z}$, $K_1 = \mathbb{Z}$, and $K_2 = \mathbb{Z} \times \mathbb{Z}$, then $G_1 = K_1 \times Q$ and $G_2 = K_2 \times Q$ are isomorphic, even though $K_1 \not\cong K_2$

Sketch of Proof

- Sketch of Proof

Sketch of Proof

- Sketch of Proof

Lemma 3

*Let $A, B, C,$ and D be nontrivial groups and let $\phi : A \times B \rightarrow C * D$ be an epimorphism. Then either $\phi(A \times \{1\})$ is all of $C * D$ and $\phi(\{1\} \times B)$ is trivial or $\phi(A \times \{1\})$ is trivial and $\phi(\{1\} \times B)$ is all of $C * D$*

Sketch of Proof

- Sketch of Proof

Lemma 3

*Let $A, B, C,$ and D be nontrivial groups and let $\phi : A \times B \rightarrow C * D$ be an epimorphism. Then either $\phi(A \times \{1\})$ is all of $C * D$ and $\phi(\{1\} \times B)$ is trivial or $\phi(A \times \{1\})$ is trivial and $\phi(\{1\} \times B)$ is all of $C * D$*

- **Remark** This is really the grain of sand that led to the pearl that is the theorem. Everything must commute and the domain, and nothing can commute in the range

Sketch of Proof

- Sketch of Proof

Lemma 3

*Let $A, B, C,$ and D be nontrivial groups and let $\phi : A \times B \rightarrow C * D$ be an epimorphism. Then either $\phi(A \times \{1\})$ is all of $C * D$ and $\phi(\{1\} \times B)$ is trivial or $\phi(A \times \{1\})$ is trivial and $\phi(\{1\} \times B)$ is all of $C * D$*

- **Remark** This is really the grain of sand that led to the pearl that is the theorem. Everything must commute and the domain, and nothing can commute in the range
- The proof uses the fact that a free product is never an internal direct product

Sketch of Proof

Lemma 4 (The Straightening-Up Lemma ($n = m$))

Let $n = m$, let K be a free product, and let $\psi : K \times K \times \dots \times K$ (n copies) $\rightarrow K \times K \times \dots \times K$ (m copies) be an isomorphism. Write $\psi_{i,j}$ for $\pi_{K_j} \circ \psi|_{K_i}$. Then ψ splits as n isomorphisms $\psi_{i,\sigma(i)}$, with σ a permutation, with all other $\psi_{i,j}$'s being the trivial map

Sketch of Proof

Lemma 4 (The Straightening-Up Lemma ($n = m$))

Let $n = m$, let K be a free product, and let $\psi : K \times K \times \dots \times K$ (n copies) $\rightarrow K \times K \times \dots \times K$ (m copies) be an isomorphism. Write $\psi_{i,j}$ for $\pi_{K_j} \circ \psi|_{K_i}$. Then ψ splits as n isomorphisms $\psi_{i,\sigma(i)}$, with σ a permutation, with all other $\psi_{i,j}$'s being the trivial map

Lemma 5 (The Straightening-Up Corollary ($n > m$))

Let $n > m$, let K be a free product of Hopfian groups, and let $\psi : K \times K \times \dots \times K$ (n copies) $\rightarrow K \times K \times \dots \times K$ (m copies) be an epimorphism. Write $\psi_{i,j}$ for $\pi_{K_j} \circ \psi|_{K_i}$. Then ψ splits as m isomorphisms $\psi_{\sigma^{-1}(i),i}$, with σ a permutation, with all other $\psi_{i,j}$'s being the trivial map

Sketch of Proof

- For the next 2 Lemmas, we have

Sketch of Proof

- For the next 2 Lemmas, we have
- K a free product of 2 copies of the same centerless, superperfect, freely indecomposable, co-Hopfian group $S_1 * S_2$ (Hopfian for the Corollary) with an unpermutable set set A (chosen as a subset of S_1)

Sketch of Proof

- For the next 2 Lemmas, we have
- K a free product of 2 copies of the same centerless, superperfect, freely indecomposable, co-Hopfian group $S_1 * S_2$ (Hopfian for the Corollary) with an unpermutable set set A (chosen as a subset of S_1)
- $G_1 = (K \times K \times \dots \times K) \rtimes_{\phi_{u_1}, \phi_{u_2}, \dots, \phi_{u_n}} \mathbb{Z}$ and

Sketch of Proof

- For the next 2 Lemmas, we have
- K a free product of 2 copies of the same centerless, superperfect, freely indecomposable, co-Hopfian group $S_1 * S_2$ (Hopfian for the Corollary) with an unpermutable set set A (chosen as a subset of S_1)
- $G_1 = (K \times K \times \dots \times K) \rtimes_{\phi_{u_1}, \phi_{u_2}, \dots, \phi_{u_n}} \mathbb{Z}$ and
- $G_2 = (S \times K \times \dots \times K) \rtimes_{\phi_{v_1}, \phi_{v_2}, \dots, \phi_{v_m}} \mathbb{Z}$ are semi-direct products

Sketch of Proof

- For the next 2 Lemmas, we have
- K a free product of 2 copies of the same centerless, superperfect, freely indecomposable, co-Hopfian group $S_1 * S_2$ (Hopfian for the Corollary) with an unpermutable set set A (chosen as a subset of S_1)
- $G_1 = (K \times K \times \dots \times K) \rtimes_{\phi_{u_1}, \phi_{u_2}, \dots, \phi_{u_n}} \mathbb{Z}$ and
- $G_2 = (S \times K \times \dots \times K) \rtimes_{\phi_{v_1}, \phi_{v_2}, \dots, \phi_{v_m}} \mathbb{Z}$ are semi-direct products
- where ϕ_{u_i} is the outer action of \mathbb{Z} on K_i given by

$$\phi_{u_i}(z)(p) = \begin{cases} p & \text{if } p \in S_1 \\ u_i^{-z} p u_i^z & \text{if } p \in S_2 \end{cases}$$

Sketch of Proof

- For the next 2 Lemmas, we have
- K a free product of 2 copies of the same centerless, superperfect, freely indecomposable, co-Hopfian group $S_1 * S_2$ (Hopfian for the Corollary) with an unpermutable set set A (chosen as a subset of S_1)
- $G_1 = (K \times K \times \dots \times K) \rtimes_{\phi_{u_1}, \phi_{u_2}, \dots, \phi_{u_n}} \mathbb{Z}$ and
- $G_2 = (S \times K \times \dots \times K) \rtimes_{\phi_{v_1}, \phi_{v_2}, \dots, \phi_{v_m}} \mathbb{Z}$ are semi-direct products
- where ϕ_{u_i} is the outer action of \mathbb{Z} on K_i given by
$$\phi_{u_i}(z)(p) = \begin{cases} p & \text{if } p \in S_1 \\ u_i^{-z} p u_i^z & \text{if } p \in S_2 \end{cases}$$
- (This particular kind of outer action is called a **partial conjugation**)

Lemma 6 (The Conder Isomorphism Lemma ($n = m$))

Let $n = m$, and let $\theta : G_1 \rightarrow G_2$ be an isomorphism. Then θ restricts to an isomorphism on the commutator subgroup $C = K \times K \times \dots \times K$, and the K factors which correspond by the Straightening-Up Lemma ($n = m$) have ϕ_{u_i} 's being determined by the same u_i in the definition of the unpermutable group A

Sketch of Proof

Lemma 6 (The Conder Isomorphism Lemma ($n = m$))

Let $n = m$, and let $\theta : G_1 \rightarrow G_2$ be an isomorphism. Then θ restricts to an isomorphism on the commutator subgroup $C = K \times K \times \dots \times K$, and the K factors which correspond by the Straightening-Up Lemma ($n = m$) have ϕ_{u_i} 's being determined by the same u_i in the definition of the unpermutable group A

Lemma 7 (The Conder Isomorphism Corollary ($n > m$))

Let $n > m$, and let $\theta : G_1 \rightarrow G_2$ be an epimorphism. Then θ restricts to an epimorphism on the commutator subgroup $C = K \times K \times \dots \times K$, and the K factors which correspond by the Straightening-Up Corollary ($n > m$) have ϕ_{u_i} 's being determined by the same u_i in the definition of the unpermutable group A

Sketch of Proof

- (\Rightarrow) Suppose there is an isomorphism θ between the two extensions. Then θ must preserve the commutator subgroup, a characteristic group, so it induces an automorphism of $K_1 \times \dots \times K_n$, say ψ . By Lemma 4, θ must send each of the n factors of $K_1 \times \dots \times K_n$ in the domain isomorphically onto exactly one of the n factors of $K_1 \times \dots \times K_n$ in the range. Let σ be the permutation from Lemma 4.

Sketch of Proof

- (\Rightarrow) Suppose there is an isomorphism θ between the two extensions. Then θ must preserve the commutator subgroup, a characteristic group, so it induces an automorphism of $K_1 \times \dots \times K_n$, say ψ . By Lemma 4, θ must send each of the n factors of $K_1 \times \dots \times K_n$ in the domain isomorphically onto exactly one of the n factors of $K_1 \times \dots \times K_n$ in the range. Let σ be the permutation from Lemma 4.
- Also, the associated map to θ on quotient groups must send the infinite cyclic quotient $G_{(k_1, \dots, k_n)} / (K_1 \times \dots \times K_n)$ isomorphically onto the infinite cyclic quotient $G_{(l_1, \dots, l_n)} / (K_1 \times \dots \times K_n)$. So, θ takes the generator, z , of \mathbb{Z} in $G_{(k_1, \dots, k_n)}$ to the an element cw^e in $G_{(l_1, \dots, l_n)}$, where c is some element of $K_1 \times \dots \times K_n$, w generates the \mathbb{Z} in $G_{(l_1, \dots, l_n)}$, and e is $+1$ or -1 .

Sketch of Proof

- But also we know that z centralises the factor $S_{i,1}$ in K_i , so its θ -image cw^e must centralize $\theta(S_{i,1}) = \psi(S_{i,1}) = Q_i$, say, in $K_{\sigma(i)}$, the copy of K to which K_i is sent under the isomorphism given by Lemma 4, and act as conjugation by $l_{\sigma(i)}$ on $\theta(S_2) = \psi(S_2) = R_i$, say, in $K_{\sigma(i)}$.

Sketch of Proof

- But also we know that z centralises the factor $S_{i,1}$ in K_i , so its θ -image cw^e must centralize $\theta(S_{i,1}) = \psi(S_{i,1}) = Q_i$, say, in $K_{\sigma(i)}$, the copy of K to which K_i is sent under the isomorphism given by Lemma 4, and act as conjugation by $l_{\sigma(i)}$ on $\theta(S_2) = \psi(S_2) = R_i$, say, in $K_{\sigma(i)}$.
- By the Kurosh Subgroup Theorem,
 $Q_i = F * B_1^{d_1} * B_2^{d_2} * \dots * B_j^{d_j}$, where F is free, each $d \in K_{\sigma(i)}$ and $B_j \leq S_{\sigma(i), u_j}$, $u_j \in \{1, 2\}$. Since the Abelianization of Q is trivial, we have that F is trivial. As $Q_i \cong S$ and S is freely indecomposable, we must have that $j = 1$. As S is co-Hopfian, we must have that $B_1 = S_{\sigma(i), u_j}$. Now, this implies that conjugation by c has the same effect as conjugation by w^{-e} on the subgroups Q_i of $K_{\sigma(i)}$, which is isomorphic to $S_{i,1}$, and Q_j of $K_{\sigma(j)}$, which is isomorphic to $S_{j,1}$, for $i \neq j$.

Sketch of Proof

- This implies $Q_i^c = Q_i^{w^{-e}}$ pointwise and $Q_j^c = Q_j^{w^{-e}}$ pointwise; as this would require $c \in K_{\sigma_i} \cap K_{\sigma_j}$ for $i \neq j$, we conclude $c = e$. Thus θ takes z to w^e .

Sketch of Proof

- This implies $Q_i^c = Q_i^{w^{-e}}$ pointwise and $Q_j^c = Q_j^{w^{-e}}$ pointwise; as this would require $c \in K_{\sigma i} \cap K_{\sigma j}$ for $i \neq j$, we conclude $c = e$. Thus θ takes z to w^e .
- So, $Q_i = \theta(S_{i,1}) = S_{\sigma(i),u_j}^d$, with $d \in K_{\sigma(i)}$ and $u_1 \in \{1, 2\}$.

Sketch of Proof

- This implies $Q_i^c = Q_i^{w^{-e}}$ pointwise and $Q_j^c = Q_j^{w^{-e}}$ pointwise; as this would require $c \in K_{\sigma_i} \cap K_{\sigma_j}$ for $i \neq j$, we conclude $c = e$. Thus θ takes z to w^e .
- So, $Q_i = \theta(S_{i,1}) = S_{\sigma(i),u_j}^d$, with $d \in K_{\sigma(i)}$ and $u_1 \in \{1, 2\}$.
- By a symmetric argument, $R = E * C_1^{f_1} * C_2^{f_2} * \dots * C_j^{f_j}$, where E is free, each $f \in S_{\sigma(i)}$ and $C_i \leq S_{\sigma(i),v_i}$, $v_i \in \{1, 2\}$ and $R = \theta(S_{i,2}) = S_{\sigma(i),v_i}^f$, with $f \in K_{\sigma(i)}$ and $v_i \in \{1, 2\}$.

Sketch of Proof

- This implies $Q_i^c = Q_i^{w^{-e}}$ pointwise and $Q_j^c = Q_j^{w^{-e}}$ pointwise; as this would require $c \in K_{\sigma_i} \cap K_{\sigma_j}$ for $i \neq j$, we conclude $c = e$. Thus θ takes z to w^e .
- So, $Q_i = \theta(S_{i,1}) = S_{\sigma(i),u_j}^d$, with $d \in K_{\sigma(i)}$ and $u_1 \in \{1, 2\}$.
- By a symmetric argument, $R = E * C_1^{f_1} * C_2^{f_2} * \dots * C_j^{f_j}$, where E is free, each $f \in S_{\sigma(i)}$ and $C_i \leq S_{\sigma(i),v_i}$, $v_i \in \{1, 2\}$ and $R = \theta(S_{i,2}) = S_{\sigma(i),v_i}^f$, with $f \in K_{\sigma(i)}$ and $v_i \in \{1, 2\}$.
- Since θ is onto, the restriction of θ to K_i must be onto. If $u_i = v_i$, then the projection onto the $S_{i,r}$ factor of $\theta|_{K_i}$, where $r \in \{1, 2\} \setminus \{u_i\}$, would be trivial, and $\theta|_{K_i}$ would not be onto. Thus, $v_i \in \{1, 2\} \setminus \{u_i\}$.

Sketch of Proof

- Suppose for some K_i , $\theta(S_{i,1}) = S_{\sigma(i),2}^d$ and $\theta(S_{i,2}) = S_{\sigma(i),1}^f$. Let $x \in S_{i,2}$ be an element other than the identity element and let $y = \theta(x) \in Q$. Then

Sketch of Proof

- Suppose for some K_i , $\theta(S_{i,1}) = S_{\sigma(i),2}^d$ and $\theta(S_{i,2}) = S_{\sigma(i),1}^f$.
Let $x \in S_{i,2}$ be an element other than the identity element and let $y = \theta(x) \in Q$. Then
- $\theta(x^z) = \theta(x)^{\theta(z)}$
 $\theta(x^{k_i}) = y^{c_{\sigma(i)} l_{\sigma(i)}^e}$
 $\theta(x^{k_i}) = y$ as conjugation by $c_{\sigma(i)} l_{\sigma(i)}$ must act trivially on Q
 $\theta(x^{k_i}) = \theta(x)$, which shows θ is not 1-1 for all $k_i \in A$ with k_i not the identity, which contradicts the fact that θ is an isomorphism.

Sketch of Proof

- Suppose for some K_i , $\theta(S_{i,1}) = S_{\sigma(i),2}^d$ and $\theta(S_{i,2}) = S_{\sigma(i),1}^f$.
Let $x \in S_{i,2}$ be an element other than the identity element and let $y = \theta(x) \in Q$. Then
- $\theta(x^z) = \theta(x)^{\theta(z)}$
 $\theta(x^{k_i}) = y^{c_{\sigma(i)} l_{\sigma(i)}^e}$
 $\theta(x^{k_i}) = y$ as conjugation by $c_{\sigma(i)} l_{\sigma(i)}$ must act trivially on Q
 $\theta(x^{k_i}) = \theta(x)$, which shows θ is not 1-1 for all $k_i \in A$ with k_i not the identity, which contradicts the fact that θ is an isomorphism.
- It follows that $\theta(S_1) = S_1^d$ and $\theta(S_2) = S_2^f$

Sketch of Proof

- Suppose for some K_i , $\theta(S_{i,1}) = S_{\sigma(i),2}^d$ and $\theta(S_{i,2}) = S_{\sigma(i),1}^f$.
Let $x \in S_{i,2}$ be an element other than the identity element and let $y = \theta(x) \in Q$. Then
- $\theta(x^z) = \theta(x)^{\theta(z)}$
 $\theta(x^{k_i}) = y^{c_{\sigma(i)} l_{\sigma(i)}^e}$
 $\theta(x^{k_i}) = y$ as conjugation by $c_{\sigma(i)} l_{\sigma(i)}$ must act trivially on Q
 $\theta(x^{k_i}) = \theta(x)$, which shows θ is not 1-1 for all $k_i \in A$ with k_i not the identity, which contradicts the fact that θ is an isomorphism.
- It follows that $\theta(S_1) = S_1^d$ and $\theta(S_2) = S_2^f$
- Also, $c_{\sigma(i)}$ is trivial, by an argument similar to the one that c is trivial.

Sketch of Proof

- Finally, if $k_i, l_{\sigma(i)} \in A$ and x is any element of the $S_{i,2}$ factor in K_i , set $y = \theta(x) = \psi(x)$. Then

$$\theta(x^z) = \theta(x^{k_i}) = \theta(x)^{\theta(k_i)} = y^{\psi(k_i)}$$

Sketch of Proof

- Finally, if $k_i, l_{\sigma(i)} \in A$ and x is any element of the $S_{i,2}$ factor in K_i , set $y = \theta(x) = \psi(x)$. Then

$$\theta(x^z) = \theta(x^{k_i}) = \theta(x)^{\theta(k_i)} = y^{\psi(k_i)}$$

- while on the other hand,

$$\theta(x^z) = \theta(x)^{\theta(z)} = y^{we} = y^{(l_{\sigma(i)})^e}$$

Sketch of Proof

- Finally, if $k_i, l_{\sigma(i)} \in A$ and x is any element of the $S_{i,2}$ factor in K_i , set $y = \theta(x) = \psi(x)$. Then

$$\theta(x^z) = \theta(x^{k_i}) = \theta(x)^{\theta(k_i)} = y^{\psi(k_i)}$$

- while on the other hand,

$$\theta(x^z) = \theta(x)^{\theta(z)} = y^{we} = y^{(l_{\sigma(i)})^e}$$

- and it follows that $y^{\psi(k_i)} = y^{(l_{\sigma(i)})^e}$, so that $\psi(k_i) = (l_{\sigma(i)})^e$; since no isomorphism of $S_{i,1}$ takes any given element of A onto any other element of A or the inverse of any other element of A , we must have $k_i = l_{\sigma(i)}^{\pm 1}$. QED Lemma

Sketch of Proof

- This is really the key lemma that changes one to allow groups with unpermutable sets instead of requiring groups with torsion elements of all orders for the kernel groups.

Sketch of Proof

- This is really the key lemma that changes one to allow groups with unpermutable sets instead of requiring groups with torsion elements of all orders for the kernel groups.
- The rest of the proof is the same as in my dissertation.

Sketch of Proof

- This is really the key lemma that changes one to allow groups with unpermutable sets instead of requiring groups with torsion elements of all orders for the kernel groups.
- The rest of the proof is the same as in my dissertation.
- In my disseratation, we had one group, Thompson's group V , that met all the requirements for a kernel group and had torsion elements of all orders.

Sketch of Proof


- This is really the key lemma that changes one to allow groups with unpermutable sets instead of requiring groups with torsion elements of all orders for the kernel groups.
- The rest of the proof is the same as in my dissertation.
- In my dissertation, we had one group, Thompson's group V , that met all the requirements for a kernel group and had torsion elements of all orders.
- Already, due to a suggestion by Jason Manning, we have a countable collection of fundamental groups of hyperbolic 3-manifolds that are allowable and meet all the requirements for a kernel group, for each of which we produce an uncountable collection of pseudo-collars.

Sketch of Proof

Now, for $\Omega = \prod_{i=1}^{\infty} \{0, 1\}$, we have the following binary tree

Sketch of Proof

Now, for $\Omega = \prod_{i=1}^{\infty} \{0, 1\}$, we have the following binary tree




Sketch of Proof

We start at the root of the tree with \mathbb{Z} , and keep blowing this quotient group up by a semi-direct product with S at each node

Sketch of Proof

We start at the root of the tree with \mathbb{Z} , and keep blowing this quotient group up by a semi-direct product with S at each node




Sketch of Proof

This gives rise to binary tree of group extensions ...

Sketch of Proof

This gives rise to binary tree of group extensions ...




Sketch of Proof

... and semi-s-cobordisms

Sketch of Proof

... and semi-s-cobordisms



Sketch of Proof


- Now, for each $\omega \in \Omega$, let (ω, n) be the corresponding finite sequence of length n

Sketch of Proof

- Now, for each $\omega \in \Omega$, let (ω, n) be the corresponding finite sequence of length n
- Corresponding to each $\omega \in \Omega$, we have an inverse sequence of groups $(G_{(\omega, n)}, \alpha_{(\omega, n)})$

Sketch of Proof

- Now, for each $\omega \in \Omega$, let (ω, n) be the corresponding finite sequence of length n
- Corresponding to each $\omega \in \Omega$, we have an inverse sequence of groups $(G_{(\omega, n)}, \alpha_{(\omega, n)})$
- Two inverse sequences are **pro-isomorphic** if and only if, after passing to subsequences, they may be put into a ladder diagram



Sketch of Proof

- To finish off the proof, let ω and η be distinct sequences in Ω

Sketch of Proof

- To finish off the proof, let ω and η be distinct sequences in Ω
- Suppose ω and η agree up to some level n_0

Sketch of Proof

- To finish off the proof, let ω and η be distinct sequences in Ω
- Suppose ω and η agree up to some level n_0
- Consider the 1-ended, pseudo-collarable $(n + 1)$ -manifolds V_ω and V_η

Sketch of Proof

- To finish off the proof, let ω and η be distinct sequences in Ω
- Suppose ω and η agree up to some level n_0
- Consider the 1-ended, pseudo-collarable $(n + 1)$ -manifolds V_ω and V_η
- Suppose, after passing to subsequences, we have their pro-fundamental group systems at infinity fitting into a ladder diagram

$$\begin{array}{ccccccc} & & \alpha & & \alpha & & \alpha \\ & & \longleftarrow & & \longleftarrow & & \longleftarrow \\ G_{(\omega, n_0)} & & G_{(\omega, n_2)} & & G_{(\omega, n_4)} & & \dots \\ & \swarrow f_{m_1} & & \swarrow f_{m_3} & & \swarrow f_{m_5} & \\ & H_{(\eta, m_1)} & & H_{(\eta, m_3)} & & H_{(\eta, m_5)} & \\ & & \beta & & \beta & & \\ & & \longleftarrow & & \longleftarrow & & \longleftarrow \\ & & & & & & \dots \end{array}$$

Lemma 8

By passing to a further subsequence if necessary, we may assume
 $n_0 < m_1 < n_2 < m_3 < \dots$

Lemma 8

By passing to a further subsequence if necessary, we may assume
 $n_0 < m_1 < n_2 < m_3 < \dots$

- Now, g_{n_2} fits the form for the Conder Isomorphism Corollary ($n > m$), so it must be onto m_1 copies of S and corresponding copies of S must have ϕ_{u_i} 's with the same u_i 's in corresponding copies of the S 's

Lemma 8

By passing to a further subsequence if necessary, we may assume $n_0 < m_1 < n_2 < m_3 < \dots$

- Now, g_{n_2} fits the form for the Conder Isomorphism Corollary ($n > m$), so it must be onto m_1 copies of S and corresponding copies of S must have ϕ_{u_i} 's with the same u_i 's in corresponding copies of the S 's
- But, ω and η only agree up to n_0 and cannot have ϕ_{u_i} 's with the same u_i 's on the remaining $m_1 - n_0$ corresponding copies of S !

Lemma 8

By passing to a further subsequence if necessary, we may assume $n_0 < m_1 < n_2 < m_3 < \dots$

- Now, g_{n_2} fits the form for the Conder Isomorphism Corollary ($n > m$), so it must be onto m_1 copies of S and corresponding copies of S must have ϕ_{u_i} 's with the same u_i 's in corresponding copies of the S 's
- But, ω and η only agree up to n_0 and cannot have ϕ_{u_i} 's with the same u_i 's on the remaining $m_1 - n_0$ corresponding copies of S !
- This concludes the proof

The End

- THE END